This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1) (Currently Amended) A compound of formula (I)

ı

or a pharmaceutically acceptable salt, or metabolite thereof, wherein

A is phenyl, naphthyl, mono- or bi-cyclic heteroaryl, or a group of the formula

optionally substituted with 1-4 substituents which are independently R^1 , OR^1 , $S(O)_pR^1$, $C(O)R^1$, $C(O)OR^1$, $C(O)NR^1R^2$, halogen, hydroxy, <u>oxide</u>, amino, cyano, or nitro;

B is phenyl, naphthyl, or pyridyl, optionally substituted with 1-4 substituents which are independently C_1 - C_5 linear or branched alkyl, C_1 - C_5 linear or branched haloalkyl, C_1 - C_3 alkoxy, hydroxy, <u>oxide</u>, amino, C_1 - C_3 alkylamino, C_1 - C_6 dialkylamino, halogen, cyano, or nitro;

L is

- (a) $-(CH_2)_m O (CH_2)_{l-1}$
- (b) $-(CH_2)_m (CH_2)_{l-1}$
- (c) $-(CH_2)_m C(O) (CH_2)_{1-}$
- (d) $-(CH_2)_m NR^3 (CH_2)_{l^-}$
- (e) $-(CH_2)_m NR^3C(O) (CH_2)_{l-1}$
- $(f) (CH_2)_m S (CH_2)_{l-1}$

(g) $-(CH_2)_m - C(O)NR^3 - (CH_2)_{l-}$, or

(h) a single bond;

m and I are integers independently selected from 0-4;

M is a pyridine ring, optionally substituted with 1-3 substituents which are independently C_1 - C_5 linear or branched alkyl, C_1 - C_5 linear or branched haloalkyl, C_1 - C_3 alkoxy, hydroxy, oxide, amino, C_1 - C_3 alkylamino, C_1 - C_6 dialkylamino, halogen, or nitro;

Q is:

(1) C(S)NR⁴R⁵;

(2) $C(O)NR^7-NR^4R^5$;

(3) tetrazolyl;

(4) imidazolyl;

(5) imidazoline-2-yl;

(6) 1,3,4-oxadiazoline-2-yl;

(7) 1,3-thiazoline-2-yl;

(8) 5-thioxo-4,5-dihydro-1,3,4-thiazoline-2-yl;

(9) 5-oxo-4,5-dihydro-1,3,4-oxadiazoline-2-yl; or

(10) a group of the formula

wherein each of R^1 , R^2 , R^3 , R^4 and R^5 is independently

(a) hydrogen,

(b) C₁-C₅ linear, branched, or cyclic alkyl,

(c) phenyl,

(d) C₁-C₃ phenyl-alkyl,

(e) up to per-halo substituted C₁-C₅ linear or branched alkyl, or

(f) - $(CH_2)_q$ -X, where X is a 5 or 6 membered heterocyclic ring, containing at least one atom selected from oxygen, nitrogen and sulfur, which is saturated, partially saturated, or aromatic, or a 8-10 membered bicyclic heteroaryl having 1-4 heteroatoms selected from the group consisting of O, N and S;

 R^4 and R^5 may additionally be taken together to form a 5 or 6 membered aliphatic ring, which may be interrupted by an atom selected from N, O or S, optionally substituted with 1-3 substituents which are independently C_1 - C_5 linear or branched alkyl, up to perhalo substituted C_1 - C_5 linear or branched alkyl, C_1 - C_3 alkoxy, hydroxy, oxo, carboxy, amino, C_1 - C_3 alkylamino, C_1 - C_6 dialkylamino, halogen, cyano, or nitro;

R⁶ is independently

- (a) hydrogen,
- (b) C₁-C₅ linear, branched, or cyclic alkyl,
- (c) cyano,
- (d) nitro,
- (e) up to per-halo substituted C₁-C₅ linear or branched alkyl. or
- (f) $-C(O)R^7$, where R^7 is C_1-C_5 linear, branched, or cyclic alkyl;

 R^7 is hydrogen or linear, branched, or cyclic $C_1\text{-}C_5$ alkyl;

q is an integer 0, 1, 2, 3, or 4 and

p is an integer 0, 1, or 2.

- 2) (Original) A compound of claim 1 wherein B is phenyl or pyridinyl, optionally substituted with 1-4 halogen.
- 3) (Original) A compound of claim 1 wherein L is -O- and B is phenyl or pyridinyl, optionally substituted with 1-4 halogen.
 - 4) (Currently Amended) A compound of claim 1 formula (I)

or a pharmaceutically acceptable salt, or metabolite thereof, wherein A is phenyl, naphthyl, indazolyl, quinolinyl, pyridyl, benzo[1,3]dioxolan-5-yl, 2,3-dihydrobenzo[1,4]dioxin-6-yl or 4H-benzo[1,3]dioxin-6-yl, optionally substituted with 1-4 substituents which are independently R¹ and halogen,

L is -O- and B is phenyl, optionally substituted with 1-4 halogen;

M is a pyridine ring, optionally substituted with 1-3 substituents which are independently C_1 - C_5 linear or branched alkyl, C_1 - C_5 linear or branched haloalkyl, C_1 - C_3 alkoxy, hydroxy, oxide, amino, C_1 - C_3 alkylamino, C_1 - C_6 dialkylamino, halogen, or nitro;

<u>Q is:</u>

- (1) C(S)NR⁴R⁵;
- (2) $C(O)NR^7 NR^4R^5$;
- (3) tetrazolyl;
- (4) imidazolyl;
- (5) imidazoline-2-yl;
- (6) 1,3,4-oxadiazoline-2-yl;
- (7) 1,3-thiazoline-2-yl;
- (8) 5-thioxo-4,5-dihydro-1,3,4-thiazoline-2-yl;
- (9) 5-oxo-4,5-dihydro-1,3,4-oxadiazoline-2-yl; or
- (10) a group of the formula

wherein each of R¹, R², R³, R⁴ and R⁵ is independently

- (a) hydrogen,
- (b) C₁-C₅ linear, branched, or cyclic alkyl,

- (c) phenyl,
- (d) C₁-C₃ phenyl-alkyl,
- (e) up to per-halo substituted C₁-C₅ linear or branched alkyl, or
- (f) -(CH₂)_g-X, where X is a 5 or 6 membered heterocyclic ring, containing at least one atom selected from oxygen, nitrogen and sulfur, which is saturated, partially saturated, or aromatic, or a 8-10 membered bicyclic heteroaryl having 1-4 heteroatoms selected from the group consisting of O, N and S;

 R^4 and R^5 may additionally be taken together to form a 5 or 6 membered aliphatic ring, which may be interrupted by an atom selected from N, O or S, optionally substituted with 1-3 substituents which are independently C_1 - C_5 linear or branched alkyl, up to perhalo substituted C_1 - C_5 linear or branched alkyl, C_1 - C_3 alkoxy, hydroxy, oxo, carboxy, amino, C_1 - C_3 alkylamino, C_1 - C_6 dialkylamino, halogen, cyano, or nitro;

R⁶ is independently

- (a) hydrogen,
- (b) C₁-C₅ linear, branched, or cyclic alkyl,
- (c) cyano,
- (d) nitro,
- (e) up to per-halo substituted C₁-C₅ linear or branched alkyl. or
- (f) -C(O)R⁷, where R⁷ is C₁-C₅ linear, branched, or cyclic alkyl;

R⁷ is hydrogen or linear, branched, or cyclic C₁-C₅ alkyl;

q is an integer 0, 1, 2, 3, or 4 and

p is an integer 0, 1, or 2.

5) **(Original)** A compound of claim 1 wherein A and B follow one of the following combinations:

A= phenyl and B= phenyl,

A= indazolyl and B= phenyl,

A= quinolinyl and B= phenyl,

A= 4H-benzo[1,3]dioxin-6-yl and B= phenyl;

A= phenyl and B= pyridyl,

A= indazolyl and B= pyridyl,

A= quinolinyl and B= pyridyl, or

A= 4H-benzo[1,3]dioxin-6-yl and B= pyridyl.

6) (Original) A compound which is

- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(hydrazinocarbonyl)pyridin-4-yl]oxy}phenyl)urea
- N-(4-{[2-(hydrazinocarbonyl)pyridin-4-yl]oxy}phenyl)-N'-(2,2,4,4-tetrafluoro-4H-1,3-benzodioxin-6-yl)urea
- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-[3-({2-[(2,2-dimethylhydrazino)carbonyl]pyridin-4-yl}oxy)phenyl]urea
- 4-{3-[({[4-chloro-3-(trifluoromethyl)phenyl]amino}carbonyl)amino]phenoxy}-N-piperidin-1-ylpyridine-2-carboxamide
- N-piperidin-1-yl-4-[3-({[(2,2,4,4-tetrafluoro-4H-1,3-benzodioxin-6-yl)amino]carbonyl}amino)phenoxy]pyridine-2-carboxamide
- 4-{3-[({[4-chloro-3-(trifluoromethyl)phenyl]amino}carbonyl)amino]phenoxy}-N-morpholin-4-ylpyridine-2-carboxamide
- N-morpholin-4-yl-4-[3-({[(2,2,4,4-tetrafluoro-4H-1,3-benzodioxin-6-yl)amino]carbonyl}amino)phenoxy]pyridine-2-carboxamide
- 4-[3-({[(1-methyl-1H-indazol-5-yl)amino]carbonyl}amino)phenoxy]-N-morpholin-4ylpyridine-2-carboxamide
- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(1H-tetrazol-5-yl)pyridin-4-yl]oxy}phenyl)urea
- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(4,5-dihydro-1H-imidazol-2-yl)pyridin-4-yl]oxy}phenyl)urea
- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(1,3,4-oxadiazol-2-yl)pyridin-4-yl]oxy}phenyl)urea
- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(4-methyl-1,3-thiazol-2-yl)pyridin-4-yl]oxy}phenyl)urea
- N-quinolin-6-yl-N'-(4-{[2-(5-thioxo-4,5-dihydro-1,3,4-thiadiazol-2-yl)pyridin-4-yl]oxy}phenyl)urea

- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(5-oxo-4,5-dihydro-1,3,4-oxadiazol-2-yl)pyridin-4-yl]oxy}phenyl)urea
- N-(4-{[2-(5-oxo-4,5-dihydro-1,3,4-oxadiazol-2-yl)pyridin-4-yl]oxy}phenyl)-N' (2,2,4,4-tetrafluoro-4H-1,3-benzodioxin-6-yl)urea
- 4-{4-[({[4-chloro-3-(trifluoromethyl)phenyl]amino}carbonyl)amino]phenoxy}-N-methylpyridine-2-carboximidamide
- 4-{4-[({[4-chloro-3trifluoromethyl)phenyl]amino}carbonyl)amino]phenoxy}pyridine-2carboximidamide
- N-methyl-4-[4-({[(2,2,4,4-tetrafluoro-4H-1,3-benzodioxin-6yl)amino]carbonyl}amino)phenoxy]pyridine-2-carboximidamide
- N-methyl-4-(4-{[(quinolin-6-ylamino)carbonyl]amino}phenoxy)pyridine-2carboximidamide
- 4-{4-[({[4-chloro-3-(trifluoromethyl)phenyl]amino}carbonyl)amino]phenoxy}pyridine-2-carbothioamide
- 4-(4-{[(quinolin-6-ylamino)carbonyl]amino}phenoxy)pyridine-2-carbothioamide or
- 4-[4-({[(1-methyl-1H-indazol-5-yl)amino]carbonyl}amino)phenoxy]pyridine-2carbothioamide
- 7) **(Original)** A pharmaceutical composition which comprises an effective amount of at least one compound of claim 1 and a physiologically acceptable carrier.
- 8) **(Withdrawn)** A method for treating or preventing a hyper-proliferative disorder in a human or other mammal comprising administering to a human or other mammal in need thereof a compound of claim 1.
- 9) **(Withdrawn)** A method for treating or preventing a hyper-proliferative disorder in a human or other mammal comprising administering to a human or other mammal in need thereof a compound of claim 1 and an additional anti-proliferative agent.
- 10) (Withdrawn) A method for treating or preventing cancer in a human or other mammal comprising administering to a human or other mammal in need

thereof a compound of claim 1 and a cytotoxic agent or cytostatic chemotherapeutic agent.

- 11) **(Withdrawn)** A method for treating or preventing a disease in a human or other mammal regulated by tyrosine kinase, associated with an aberration in the tyrosine kinase signal transduction pathway, comprising administering to a human or other mammal in need thereof a compound of claim 1.
- 12) **(Withdrawn)** A method for treating or preventing a disease in a human or other mammal mediated by the VEGF-induced signal transduction pathway, comprising administering to a human or other mammal in need thereof a compound of claim 1.
- 13) **(Withdrawn)** A method for treating or preventing a disease in a human or other mammal characterized by abnormal angiogenesis or hyperpermeability processes, comprising administering to a human or other mammal in need thereof a compound of claim 1.
- 14) **(Withdrawn)** A method for treating or preventing a disease in a human or other mammal characterized by abnormal angiogenesis or hyperpermeability processes, comprising administering to a human or other mammal in need thereof a compound of claim 1 simultaneously with another angiogenesis inhibiting agent in the same formulation or in separate formulations.
- 15) (Withdrawn) A method for treating or preventing one or more of the following conditions in humans and/or other mammals: tumor growth, retinopathy, ischemic retinal-vein occlusion, retinopathy of prematurity, age related macular degeneration; rheumatoid arthritis, psoriasis, a bolos disorder associated with subepidermal blister formation, including bullous pemphigoid, erythema multiforme, or dermatitis herpetiformis, comprising administering to a human or other mammal in need thereof a compound of claim 1.

(Withdrawn) A method for treating or preventing one or more of the following conditions in humans and/or other mammals: tumor growth, retinopathy, diabetic retinopathy, ischemic retinal-vein occlusion, retinopathy of prematurity, age related macular degeneration; rheumatoid arthritis, psoriasis, bullous disorder associated with subepidermal blister formation, bullous pemphigoid, erythema multiforme, and dermatitis herpetiformis, in combination with an infectious disease selected from the group consisting of: tuberculosis, Helicobacter pylori infection during peptic ulcer disease, Chaga's disease resulting from Trypanosoma cruzi infection, effects of Shiga-like toxin resulting from E. coli infection, effects of enterotoxin A resulting from Staphylococcus infection, meningococcal infection, and infections from Borrelia burgdorferi, Treponema pallidum, cytomegalovirus, influenza virus, Theiler's encephalomyelitis virus, and the human immunodeficiency virus (HIV),

said method comprising administering to a human or other mammal in need thereof a compound of claim 1.

- 17) **(Withdrawn)** A method for treating or preventing diseases mediated by the VEGF-induced signal transduction pathway comprising administering a compound selected from the group consisting of:
- 4-{4-[3-(4-Chloro-3-trifluoromethyl-phenyl)-ureido}-phenoxy}-pyridine-2carbothioic acid amide;
- 4-{3-[3-(2,2,4,4-Tetrafluoro-4H-benzo[1,3]dioxin-6-yl)-ureido]-phenoxy}-pyridine-2-carboxylic acid (1-piperidyl)-amide;
- 4-{3-[3-(2,2,4,4-Tetrafluoro-4H-benzo[1,3]dioxin-6-yl)-ureido]-phenoxy}-pyridine-2-carboxylic acid (4-morpholino)-amide;
- 4-{3-[3-(1-Methyl-1H-indazol-5-yl)-ureido]-phenoxy}-pyridine-2-carboxylic acid (4-morpholino)-amide;
- 4-{4-[3-(4-Chloro-3-trifluoromethyl-phenyl)-ureido]-phenoxy}-pyridine-2carboxamidine:
- 1-(4-Chloro-3-trifluoromethyl-phenyl)-3-{4-[2-(1H-tetrazol-5-yl)-pyridinyl-4-oxy]-phenyl}-urea;
- 1-(4-Chloro-3-trifluoromethyl-phenyl)-3-{4-[2-(4,5-dihydro-1H-imidazol-2-yl)-pyridinyl-4-oxy]-phenyl}-urea;

 4-{4-[3-(4-Chloro-3-trifluoromethyl-phenyl)-ureido]-phenoxy}-N-methyl-pyridine-2carboxamidine;

or a salt form, prodrug or metabolite thereof.

- 18) **(Withdrawn)** A method for treating or preventing cancer comprising administering a compound selected from the group consisting of:
- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(hydrazinocarbonyl)pyridin-4-yl]oxy}phenyl)urea
- N-(4-{[2-(hydrazinocarbonyl)pyridin-4-yl]oxy}phenyl)-N'-(2,2,4,4-tetrafluoro-4H-1,3-benzodioxin-6-yl)urea
- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-[3-({2-[(2,2-dimethylhydrazino)carbonyl]pyridin-4-yl}oxy)phenyl]urea
- 4-{3-[({[4-chloro-3-(trifluoromethyl)phenyl]amino}carbonyl)amino]phenoxy}-N-piperidin-1-ylpyridine-2-carboxamide
- N-piperidin-1-yl-4-[3-({[(2,2,4,4-tetrafluoro-4H-1,3-benzodioxin-6-yl)amino]carbonyl}amino)phenoxy]pyridine-2-carboxamide
- 4-{3-[({[4-chloro-3-(trifluoromethyl)phenyl]amino}carbonyl)amino]phenoxy}-N-morpholin-4-ylpyridine-2-carboxamide
- N-morpholin-4-yl-4-[3-({[(2,2,4,4-tetrafluoro-4H-1,3-benzodioxin-6-yl)amino]carbonyl}amino)phenoxy]pyridine-2-carboxamide
- 4-[3-({[(1-methyl-1H-indazol-5-yl)amino]carbonyl}amino)phenoxy]-N-morpholin-4-ylpyridine-2-carboxamide
- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(1H-tetrazol-5-yl)pyridin-4-yl]oxy}phenyl)urea
- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(4,5-dihydro-1H-imidazol-2-yl)pyridin-4-yl]oxy}phenyl)urea
- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(1,3,4-oxadiazol-2-yl)pyridin-4-yl]oxy}phenyl)urea
- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(4-methyl-1,3-thiazol-2-yl)pyridin-4-yl]oxy}phenyl)urea
- N-quinolin-6-yl-N'-(4-{[2-(5-thioxo-4,5-dihydro-1,3,4-thiadiazol-2-yl)pyridin-4-yl]oxy}phenyl)urea

- N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(4-{[2-(5-oxo-4,5-dihydro-1,3,4-oxadiazol-2-yl)pyridin-4-yl]oxy}phenyl)urea
- N-(4-{[2-(5-oxo-4,5-dihydro-1,3,4-oxadiazol-2-yl)pyridin-4-yl]oxy}phenyl)-N' (2,2,4,4-tetrafluoro-4H-1,3-benzodioxin-6-yl)urea
- 4-{4-[({[4-chloro-3-(trifluoromethyl)phenyl]amino}carbonyl)amino]phenoxy}-N-methylpyridine-2-carboximidamide
- 4-{4-[({[4-chloro-3-(trifluoromethyl)phenyl]amino}carbonyl)amino]phenoxy}pyridine-2carboximidamide
- N-methyl-4-[4-({[(2,2,4,4-tetrafluoro-4H-1,3-benzodioxin-6-yl)amino]carbonyl}amino)phenoxy]pyridine-2-carboximidamide
- N-methyl-4-(4-{[(quinolin-6-ylamino)carbonyl]amino}phenoxy)pyridine-2carboximidamide
- 4-{4-[({[4-chloro-3-(trifluoromethyl)phenyl}amino}carbonyl)amino]phenoxy}pyridine-2-carbothioamide
- 4-(4-{[(quinolin-6-ylamino)carbonyl]amino}phenoxy)pyridine-2-carbothioamide
- 4-[4-({[(1-methyl-1H-indazol-5-yl)amino]carbonyl}amino)phenoxy]pyridine-2-carbothioamide, or a salt form, prodrug or metabolite thereof.

19) (Currently Amended) A compound of formula (I)

or a pharmaceutically acceptable salt or metabolite thereof, wherein

A is

wherein A is optionally substituted with 1-4 substituents which are independently R^1 , OR^1 , $S(O)_pR^1$, $C(O)R^1$, $C(O)OR^1$, $C(O)NR^1R^2$, halogen, hydroxy, <u>oxide</u>, amino, cyano, or nitro;

B is phenyl, naphthyl, or pyridyl, optionally substituted with 1-4 substituents which are independently C_1 - C_5 linear or branched alkyl, C_1 - C_5 linear or branched haloalkyl, C_1 - C_3 alkoxy, hydroxy, <u>oxide</u>, amino, C_1 - C_3 alkylamino, C_1 - C_6 dialkylamino, halogen, cyano, or nitro;

L is

- (a) $-(CH_2)_m O (CH_2)_{l-1}$
- (b) $-(CH_2)_m-(CH_2)_{l-1}$
- (c) $-(CH_2)_m C(O) (CH_2)_{l-}$
- (d) $-(CH_2)_m NR^3 (CH_2)_{l-1}$
- (e) $-(CH_2)_{m}$ NR³C(O)-(CH₂)_i-,
- $(f) (CH_2)_m S (CH_2)_{l-1}$
- (g) $-(CH_2)_m-C(O)NR^3-(CH_2)_{l^-}$, or
- (h) a single bond;

m and I are integers independently selected from 0-4;

M is a pyridine ring, optionally substituted with 1-3 substituents which are independently C_1 - C_5 linear or branched alkyl, C_1 - C_5 linear or branched haloalkyl, C_1 - C_3 alkoxy, hydroxy, oxide, amino, C_1 - C_3 alkylamino, C_1 - C_6 dialkylamino, halogen, or nitro;

Q is:

- (1) C(S)NR⁴R⁵;
- (2) $C(O)NR^7 NR^4R^5$;
- (3) tetrazolyl;
- (4) imidazolyl;
- (5) imidazoline-2-yl;
- (6) 1,3,4-oxadiazoline-2-yl;
- (7) 1,3-thiazoline-2-yl;

- (8) 5-thioxo-4,5-dihydro-1,3,4-thiazoline-2-yl;
- (9) 5-oxo-4,5-dihydro-1,3,4-oxadiazoline-2-yl; or
- (10) a group of the formula

wherein each of R¹, R², R³, R⁴ and R⁵ is independently

- (a) hydrogen,
- (b) C₁-C₅ linear, branched, or cyclic alkyl,
- (c) phenyl,
- (d) C₁-C₃ phenyl-alkyl,
- (e) up to per-halo substituted C₁-C₅ linear or branched alkyl, or
- (f) -(CH₂)_q-X, where X is a 5 or 6 membered heterocyclic ring, containing at least one atom selected from oxygen, nitrogen and sulfur, which is saturated, partially saturated, or aromatic, or a 8-10 membered bicyclic heteroaryl having 1-4 heteroatoms selected from the group consisting of O, N and S;

 R^4 and R^5 may additionally be taken together to form a 5 or 6 membered aliphatic ring, which may be interrupted by an atom selected from N, O or S, optionally substituted with 1-3 substituents which are independently C_1 - C_5 linear or branched alkyl, up to perhalo substituted C_1 - C_5 linear or branched alkyl, C_1 - C_3 alkoxy, hydroxy, oxo, carboxy, amino, C_1 - C_3 alkylamino, C_1 - C_6 dialkylamino, halogen, cyano, or nitro;

R⁶ is independently

- (a) hydrogen,
- (b) C₁-C₅ linear, branched, or cyclic alkyl,
- (c) cyano,
- (d) nitro,
- (e) up to per-halo substituted C₁-C₅ linear or branched alkyl. or
- (f) $-C(O)R^7$, where R^7 is C_1-C_5 linear, branched, or cyclic alkyl;

R⁷ is hydrogen or linear, branched, or cyclic C₁-C₅ alkyl;

q is an integer 0, 1, 2, 3, or 4 and

p is an integer 0, 1, or 2.

- 20) **(Original)** A compound of claim 19 wherein B is phenyl or pyridinyl, optionally substituted with 1-4 halogen.
- 21) **(Original)** A compound of claim 19 wherein L is –O- and B is phenyl or pyridinyl, optionally substituted with 1-4 halogen.
- 22) (Original) A compound as in claim 19 wherein B is phenyl or pyridyl, L is -O-,

M a pyridine ring substituted only by Q, and Q is

C(S)NR⁴R⁵;

C(O)NR⁷-NR⁴R⁵;

or

a group of the formula

with each of R⁴ and R⁵, independently:

- (a) hydrogen,
- (b) C₁-C₅ linear, branched, or cyclic alkyl,
- (c) phenyl,
- (d) C₁-C₃ phenyl-alkyl,
- (e) up to per-halo substituted C₁-C₅ linear or branched alkyl, or
- (f) $-(CH_2)_q-X$, where the substituent X is pyridinyl and the variable q is preferably an integer 0 or 1, and


R⁶ is:

- (a) hydrogen,
- (b) C₁-C₅ linear, branched, or cyclic alkyl, or
- (c) cyano.

23) (Currently Amended) A compound of formula (I)

or a pharmaceutically acceptable salt or metabolite thereof, wherein

A is



wherein A is optionally substituted with 1-4 substituents which are independently R¹, OR¹, or halogen;

١

B is phenyl or pyridinyl, optionally substituted with 1-4 substituents which are independently C_1 - C_5 linear or branched alkyl, C_1 - C_5 linear or branched haloalkyl, C_1 - C_3 alkoxy, hydroxy, <u>oxide</u>, amino, C_1 - C_3 alkylamino, C_1 - C_6 dialkylamino, halogen, cyano, or nitro,

L is -O-,

M is a pyridine ring,

Q is:

- (1) C(S)NR⁴R⁵;
- (2) $C(O)NR^7 NR^4R^5$;
- (3) tetrazolyl;

- (4) imidazolyl;
- (5) imidazoline-2-yl;
- (6) 1,3,4-oxadiazoline-2-yl;
- (7) 1,3-thiazoline-2-yl;
- (8) 5-thioxo-4,5-dihydro-1,3,4-thiazoline-2-yl;
- (9) 5-oxo-4,5-dihydro-1,3,4-oxadiazoline-2-yl; or
- (10) a group of the formula

wherein each of R1, R4 and R5 is independently

- (a) hydrogen,
- (b) C₁-C₅ linear, branched, or cyclic alkyl,
- (c) phenyl,
- (d) C₁-C₃ phenyl-alkyl,
- (e) up to per–halo substituted C₁-C₅ linear or branched alkyl, or
- (f) -(CH₂)_q-X, where X is a 5 or 6 membered heterocyclic ring, containing at least one atom selected from oxygen, nitrogen and sulfur, which is saturated, partially saturated, or aromatic, or a 8-10 membered bicyclic heteroaryl having 1-4 heteroatoms selected from the group consisting of O, N and S;

 R^4 and R^5 may additionally be taken together to form a 5 or 6 membered aliphatic ring, which may be interrupted by an atom selected from N, O or S, optionally substituted with 1-3 substituents which are independently C_1 - C_5 linear or branched alkyl, up to perhalo substituted C_1 - C_5 linear or branched alkyl, C_1 - C_3 alkoxy, hydroxy, oxo, carboxy, amino, C_1 - C_3 alkylamino, C_1 - C_6 dialkylamino, halogen, cyano, or nitro;

R⁶ is independently

- (a) hydrogen,
- (b) C₁-C₅ linear, branched, or cyclic alkyl,
- (c) cyano,

- (d) nitro,
- (e) up to per-halo substituted C₁-C₅ linear or branched alkyl. or
- (f) $-C(O)R^7$, where R^7 is C_1-C_5 linear, branched, or cyclic alkyl;

 R^7 is hydrogen or linear, branched, or cyclic C_1 - C_5 alkyl; q is an integer 0, 1, 2, 3, or 4 and p is an integer 0, 1, or 2.

- 24) **(Original)** A compound of claim 23 wherein B is phenyl or pyridinyl, substituted with 1-4 halogen.
 - 25) (Original) A compound as in claim 23 wherein

M a pyridine ring substituted only by Q, and Q is

C(S)NR⁴R⁵;

C(O)NR⁷-NR⁴R⁵;

or

a group of the formula

with each of R⁴ and R⁵, independently:

- (a) hydrogen,
- (b) C₁-C₅ linear, branched, or cyclic alkyl,
- (c) phenyl,
- (d) C₁-C₃ phenyl-alkyl,
- (e) up to per-halo substituted C₁-C₅ linear or branched alkyl, or
- (f) $-(CH_2)_q-X$, where the substituent X is pyridinyl and the variable q is preferably an integer 0 or 1, and

R⁶ is:

(a) hydrogen,

- - 27) (Canceled)
 - 28) (Canceled)
- 29) (**Previously presented**) An ester derivative of a compound of formula I of claim 1.
- 30) **(Previously presented)** An ester derivative of a compound of formula I of claim 10.